

Effects of Medicare Coverage for the Chronically Ill on Health Insurance, Utilization, and Mortality: Evidence from Coverage Expansions Affecting People with End-Stage Renal Disease*

Martin S Andersen[†]

Current Draft: March 5, 2018

First Draft: August 15, 2016

Abstract

I study the effect of the 1973 expansions of Medicare coverage among individuals with end-stage renal disease (ESRD) on insurance coverage, health care utilization, and mortality. I find that the expansions increased insurance coverage by between 22 and 30 percentage points, in models that include trends in age, with the increase explained by Medicare coverage, and increased physician visits by 18 to 35 percent. These expansions also decreased mortality due to kidney disease in the under 65 population by between 0.5 and 1.0 deaths per 100,000. Lastly, I provide evidence for two mechanisms that affected mortality: an increase in access to and use of treatment, which may be due to changes in insurance coverage; and an increase in entry of dialysis clinics. In a simulation, the program saved between 2000 and 14000 life years per year, which is insufficient to offset the cost of the program.

Keywords: Insurance, Mortality, Kidney disease, Health, Health Insurance

JEL Codes: I13, I18, H51

*Aaron Ladd, James Frizzell, and Mohamad Nouredine provided excellent research assistance. Dr. Glenn Gee provided helpful insight into the coding and epidemiology of kidney disease, but, as usual, any errors or omissions are my own. I thank participants at the UNCG Brown Bag and the Southern Economic Association, American Society of Health Economists, and AEA/ASSA Annual Meetings for helpful comments and the Editor, Kitt Carpenter, and three anonymous referees for exceptionally helpful comments. As usual, any errors and omissions are my own.

[†]Department of Economics, UNC Greensboro. 516 Stirling Street, Greensboro, NC 27412 USA.

1 Introduction

The United States has typically expanded public insurance programs by providing coverage to distinct demographic groups. For example, the introduction of Medicare and Medicaid in 1966 provided insurance coverage to people who were 65 and older or had low income. However, several expansions of these programs have defined eligibility based in part on the presence of medical conditions (e.g. long-term disabled, people with end-stage renal disease, pregnant women, and women diagnosed with breast or cervical cancer). By selecting on ill-health, the effects of a disease-specific insurance expansion on insurance coverage, health care utilization, and health outcomes may differ considerably from the effects of more broad-based expansions.

Previous studies of the Medicare and Medicaid programs have demonstrated that Medicare may reduce mortality (Card et al., 2009; Chay et al., 2017), increase health care utilization (Card et al., 2008), and improve financial risk protection (Barcellos and Jacobson, 2015; Engelhardt and Gruber, 2011), while the introduction of state Medicaid programs reduced infant mortality (Goodman-Bacon, 2017). More recent evidence from an expansion of Medicaid for pregnant women demonstrates improvements in infant health outcomes (Currie and Gruber, 1996b) and a related expansion affecting children improved their health and increased health care utilization (Currie and Gruber, 1996a). A recent randomized study of the Oregon Medicaid program (Finkelstein et al., 2012) also demonstrated greater health care utilization and better self-rated physical and mental health among people randomized to receive Medicaid coverage, although there was no statistically significant difference in mortality.

There is, to my knowledge, no empirical evidence of the effects of three other disease-specific insurance expansions that provided insurance coverage for women with breast or cervical cancer, the long-term disabled, and people with end-stage renal disease (ESRD). In this paper, I examine the effect of a 1973 Medicare expansion that provided coverage to two groups of people: long-term beneficiaries of the Social Security Disability Insurance (SSDI) program and people who are undergoing dialysis or have received a kidney transplant due to having end-stage renal disease. The focus of this paper is the effect of the expansion on people with kidney disease who became eligible for Medicare coverage through either the ESRD route, if they were not already receiving SSDI payments, or due to SSDI receipt. Consistent with prior practice of the Medicare program itself, I consider both sets of enrollees as being enrolled in the ESRD program.¹

These expansions are attractive to study for several reasons. First, the introduction of the program was, for the most part, unanticipated so that there is unlikely

¹The Medicare Trustees' Reports from this period all pool the ESRD population and the disabled with ESRD populations because the disabled population with ESRD is more similar to the non-disabled ESRD population than the rest of the disabled population.

to be any significant anticipatory effects (Ball, 1973). Second, people with kidney disease tend to be in extremely poor health, so insurance is likely to have unusually large effects on health. Third, because for most people treatment was unaffordable prior to the expansion and insurance typically did not cover treatment (Congressional Research Service, 1971; Rettig, 2011), these results provide some insight into the welfare consequences of moral hazard induced spending since the bulk of any increase in utilization can be attributed to ex-post moral hazard. The ESRD program is also worthy of study on the basis of the size of the program. In 2015 the United States spent over \$30 billion to treat 500,000 Medicare beneficiaries with ESRD, which represents 1% of all Medicare beneficiaries and 5% of Medicare spending. In other words, the ESRD program is almost as large as the entire Medicaid program in the state of Texas, which is the third largest Medicaid program (by spending) in the country.

In order to identify the effect of the ESRD program, I estimate triple-difference models that compare outcomes for people over 65, who were always eligible for Medicare coverage, versus those under 65, before versus after the expansion took effect, with versus without ESRD. However, due to the expansion of Medicare coverage to the long-term disabled, the triple difference estimate is biased. Hence, I also estimate difference-in-differences models that condition on having ESRD, which yields unbiased estimates as long as there is no selection into treatment, i.e. as long as ex-ante moral hazard is small. These two estimators will yield similar results as long as either the effect of Medicare eligibility is small in the non-ESRD group or the share of people eligible for Medicare coverage in that group is small.

In this paper I document three main facts about the ESRD program. First, I demonstrate that the ESRD expansion significantly increased insurance coverage among people under 65 years of age with kidney disease. Close to the traditional Medicare eligibility threshold of 65, I find a 22.6 to 29.6 percentage point increase in the probability of any insurance coverage among people with kidney disease. I find somewhat larger increases in Medicare coverage (26.0 and 33.9 percentage points, respectively), indicating that some people would have had insurance coverage in the absence of the expansion.

Second, I find that the ESRD expansion increased physician visits by 25 to 35 percent for people with kidney disease below 65 years of age. The increase in physician visits is consistent with my results on health insurance coverage and implies that a ten percent increase in the share of the population with insurance increases physician visits by about eight percent. Because of the wording of the survey question that I use to assess physician visits, it is also possible that the increase in physician visits represents an increase in visits to, among other things, dialysis clinics.

Third, I document a significant reduction in mortality due to kidney disease of between two and seven log points, depending on specification and definition of kidney disease. I am able to replicate this finding in cross-national comparisons as well that

allow me to control for innovations in kidney disease treatment across countries. My results imply that the program averted between 174 and 325 deaths per year for whites between 45 and 64 years of age (my estimation sample). Assuming that the entire change in mortality arose among people who gained insurance coverage, then my mortality and insurance results imply that the probability of dying in the coming year of kidney disease fell by 0.2 to 0.5 percentage points.

I am also able to provide evidence in support of two mechanisms by which the ESRD expansion affected health. First, the state-specific effect of the ESRD expansion on kidney disease mortality was larger in states that had more treatment facilities per capita in 1971. One interpretation of this result is that the presence of treatment facilities reduced mortality by increasing access to treatment. This interpretation is also consistent with the increase in physician visits. Second, I document an increase in the number of dialysis clinics per capita from 1971 to 1975 in states that had a higher under 65 mortality rate due to kidney disease, which is consistent with a demand side shock encouraging entry of new treatment facilities.

My mortality estimates also allow me to extrapolate to changes in survival and imply that the expansion saved between 2000 and 14000 life years, based on the change in survival for 45 year olds. This range encompasses some values where, using a value of \$100,000 per statistical life year, the cost of the program are outweighed by the survival benefits. However, these estimates ignore other costs that the program imposes on society (e.g. increased disability insurance payments) but also ignores the value of spillover effects on to people 65 and older.

The remainder of the paper proceeds as follows. Section 2 provides background information on end-stage renal disease, discusses the role that the federal government has played in the treatment of ESRD, and describes the 1973 Medicare expansions that I study. Section 3 describes the data that I use for my analyses and the empirical approach that I take, while sections 4 present my main results from the Medicare expansion in 1973. Section 5 presents potential mechanisms behind my results. Section 6 discusses the welfare implications of my results. Section 7 concludes.

2 Background

End-stage renal disease (ESRD) is the end result of a progressive decline in kidney function due to chronic kidney disease. Leading causes of ESRD and chronic kidney disease include chronic kidney disease include diabetes, hypertension, glomerulonephritis, polycystic kidney disease, kidney stones, urinary tract infections, and various congenital defects (National Kidney Foundation, 2009).² The loss of kidney

²Appendix table A lists the ICDA-8 codes that I use to identify deaths with these underlying cause of death codes. In analyses using the National Health Interview Survey, I also include data that uses ICD-7 and ICD-9 codes, which are also identified in the appendix table.

function that characterizes ESRD leads to a rapid buildup in toxins in the blood that, left unchecked, rapidly leads to death.

Treatment for ESRD emphasizes restoring or augmenting the body's ability to filter out toxins either by transplanting a functioning kidney from either a living or cadaveric donor or by externally filtering blood using a dialysis machine. There were significant scientific advances affecting both forms of treatment in the late 1950s through the 1960s. The first successful kidney transplant was performed in 1956 with the subsequent decade leading to slow, but steady, improvements in transplantation (Congressional Research Service, 1971) so that by 1971 there were 1172 kidney transplants performed (Rettig, 1976). Throughout this period, kidney transplantation was a costly procedure with the Congressional Research Service (1971) estimated that kidney transplantation had a nominal one-time cost of \$10,000 to \$20,000 (\$59,000 to \$117,000 in 2015) and maintenance costs of \$1,000 per year (\$5,900 in 2015).

Chronic dialysis, which is what is necessary to treat ESRD, was not feasible until 1960. Furthermore, at its inception, dialysis was extremely costly leading to rationing at the first dialysis clinic in the United States (Alexander, 1962). In July of 1972, there were 5786 living dialysis patients in the United States (Rettig, 1976, p. 200) and the Congressional Research Service (1971) estimated that the annual cost of dialysis was \$15,000 in 1971 (nominal dollars, \$85,000 in 2015 using the CPI-U).

During the 1960s, the federal government took an active role in promoting the diffusion of treatments for ESRD as well as funding research and development of new treatments. In 1963, the Veteran's Administration began to open dialysis clinics in its hospitals across the country and, by 1971, there were 40 dialysis facilities and 15 transplant programs open at VA and military hospitals across the U.S. Beginning in 1964, the National Institutes of Health started programs to study transplant immunology, which was intended to increase the number of successful kidney transplants. In 1965, the Public Health Service started the Kidney Disease Control Program, which provided start-up grants to open a dozen dialysis centers (Rettig, 1991). The federal government, through the Bureau of the Budget, also began examining the fiscal implications of the growth in ESRD and the advent of new methods to treat ESRD, although these discussions ultimately did not appear to have affected federal policy (for further discussion see Rettig, 1991).

In 1972 Congress, for the first time, passed a law expanding eligibility for Medicare coverage, with the expansion taking effect on July 1, 1973. Congress did so by declaring that two groups were eligible for coverage people who : had been eligible for Social Security Disability Insurance (SSDI) benefits for more than 24 months; or have received three, or more, months of renal dialysis with coverage extending up to twelve months after a person received a kidney transplant.

Neither component of the expansion was truly universal since in both cases, only individuals who were eligible for insurance under the Social Security program became eligible. Collectively, these two programs increased Medicare enrollment by

1.7 million people, of whom 6,371 were eligible solely due to the ESRD in the first year of the program. By 1978, there were almost 44,000 Medicare beneficiaries with ESRD, of whom almost 35,000 were under 65 years of age, with per capita spending of almost \$65,000 (in 2015 dollars).

The ESRD component of the expansion (which includes long-term disabled with ESRD), which was initially expected to enroll 35,000 people and cost \$1 billion (nominal) per year, rapidly ballooned in size, covering more than 50,000 people and costing over \$1 billion per year in 1979 (table 1). In 2013, the ESRD program covered almost half a million people at a cost of \$30 billion, which represents approximately 1% of Medicare enrollees and 5% of Medicare spending.

[Table 1 about here.]

3 Data and Empirical Framework

3.1 Data

I use data from a variety of sources to measure insurance coverage, health care utilization, and mortality in my main analyses as well as data on potential mechanisms and confounding factors. In this subsection, I describe each of these data sources.

3.1.1 Insurance Coverage and Health Care Utilization

The National Health Interview Survey asked respondents about insurance coverage in even numbered years beginning in 1968, although the specific wording and universe for various questions has changed over time. In 1968 the NHIS inquired about health insurance generically and did not differentiate between public and private coverage and it was not until 1978 that the NHIS inquired about Medicare coverage for people under 65 years of age. In the 1974 and 1976 waves of the survey individuals with only Medicare coverage were instructed to respond that they were uninsured. As a result, I present results using data from 1968, 1970, 1972, 1978, and 1980 for most insurance outcomes (I include data on private insurance coverage in 1974 and 1976). I define an individual as having private insurance coverage based on whether or not an individual reported having private hospital coverage (as in Finkelstein, 2007) and define Medicare coverage in a comparable manner.

The NHIS also included questions on the number of doctor visits in the prior year beginning in 1969, which I use to measure health care utilization. Because the NHIS questions refer to treatment received over the prior year, I omit people 65 years of age from the utilization analysis and all data from July 1973 through June 1974, the twelve months following the implementation of the Medicare expansion.

I use the condition inventory and the list of conditions that caused the interviewee to miss days from work or access health care services to construct indicators for the presence of kidney disease. The coding is based on the codes for the broad

definition, but incorporating the NHIS omissions, listed in Appendix Table A. In total, out of 371,181 people in the NHIS, I identified 1644 people between 45 and 84 years of age with kidney disease using the broad definition. Despite the small sample size, the ESRD expansion is likely to have led to large changes in insurance coverage, hence I remain sufficiently powered to identify effects of the ESRD expansion on insurance coverage. For the utilization analyses, it is possible that I will be underpowered to detect effects if the increase in physician visits from the expansion is small.

3.1.2 Mortality

I use the Multiple Cause of Death files from the National Center for Health Statistics' (NCHS) for the years from 1968 through 1978 (United States Department of Health and Human Services. National Center for Health Statistics, 2007a,b). These data provide the state and county of residence, race, gender, age, underlying cause of death and all other diagnoses listed on the death certificate for all deaths in the United States, except in 1972, when the NCHS was only able to process half of the submitted death certificates.³ Preliminary analyses of the distribution of deaths by age indicated significant excess mass at five-year intervals of age for non-white individuals, which was also reported in Honoré and Lleras-Muney (2006), so I omit non-whites from my mortality analyses. I also drop deaths to non-U.S. residents since they were not eligible for the ESRD program.

I code each death as being a kidney disease death, or not, based on either the underlying cause of death, which the World Health Organization defines as “the disease or injury that initiated the train of events leading directly to death, or the circumstances of the accident or violence which produced the fatal injury,” or using any of the diagnosis codes listed on the death certificate. For each source of cause of death diagnosis codes, I defined a death as due to kidney disease using three sets of diagnosis codes. First, I defined a “narrow” definition of kidney disease, which did not restrict to only chronic disease, but is generally based on the “renal failure” codes in the ICDA-8. Second, I created a “chronic” definition by restricting the narrow definition to deaths due to chronic causes. Lastly, I created a “broad” definition, which was based on the codes used by the Kidney Disease Program in tracking kidney disease mortality (Kidney Disease Program, 1971). Appendix Table A lists the ICDA-8 diagnosis codes for the three cause of death groupings that I use.

I combine the mortality data with population data from the SEER program and the U.S. Census Bureau in order to adjust for changes in the size of the population over time, which also affects the expected number of deaths due to kidney disease. Because these data do not break out population figures for individuals 85 and over, I restrict my analysis to deaths to individuals who are 84 or younger.

³Since I use functions of the count of deaths in a given demographic-time cell as my dependent variable, I multiply the count of deaths in 1972 by 2.

3.1.3 Mechanisms and Confounders

In my discussion of mechanisms and potential confounders, below, I rely on data from three other datasets. I collected data on the geographic distribution of treatment facilities in 1971 from the publication “Kidney disease services, facilities, and programs in the United States” (Kidney Disease Program, 1971), which lists treatment facilities by state. Based on the name of the facility, I also classified these facilities into Veteran’s Administration/Military vs. civilian categories since access to the former may be restricted. Data on treatment facilities in 1975 came from the 1977 Annual Statistical Supplement to the Social Security Bulletin, which lists the number of hospital transplant programs, hospital-based dialysis clinics, and free-standing dialysis clinics by state.

I collected data on the share of people in an age-gender-state cell who receive income from either Social Security or the Supplemental Security Income program from the March CPS supplements for 1977 to 1979 (spanning 1976 to 1978).

3.2 Empirical Approach

3.2.1 Identification

My data includes three sources of variation that I could use to identify the effect of the Medicare ESRD program on insurance coverage, health care utilization, and kidney disease mortality. First, there are differences over time in Medicare eligibility for individuals of the same age and disease status. Second, there are differences by age in eligibility for Medicare for individuals in the same year and disease status. Third, there are differences by disease status in eligibility for Medicare coverage for individuals in the same year and of the same age. In principle, these three sources of variation would justify a triple difference estimator assuming that potential outcomes between these groups satisfy a “parallel trends” assumption (Lee and Kang, 2006). However, in my setting the parallel trends assumption is unlikely to hold because the SSDI expansion means that there is partial takeup of Medicare coverage in one of the comparison groups. The structure of the problem, allows me to identify the source of any bias from these comparisons and identify a solution that leads to unbiased estimates of the intent-to-treat effect of the Medicare expansion on people with kidney disease.

To demonstrate the bias and identify situations in which it does not affect my results, let Y_{akt}^e denote the potential outcome for someone in age group a ($a = 1$ for people under 65) who has kidney disease if $k = 1$, in time period t ($t = 1$ in the post period), and is either eligible ($e = 1$) or ineligible ($e = 0$) for Medicare coverage. Assume that there is a probability α_{akt} that a person is eligible for Medicare in each akt cell and define $Y_{akt} = \alpha_{akt}Y_{akt}^1 + (1 - \alpha_{akt})Y_{akt}^0$. Ignoring the fact that some people 65 and older are not eligible for Medicare, Medicare program rules imply that $\alpha_{0kt} = 1$ for all $k, t \in \{0, 1\}$ and $\alpha_{1k0} = 0$ for $k \in \{0, 1\}$. Finally, because (almost) everyone with kidney disease is automatically eligible for Medicare coverage, but

only some people without kidney disease are eligible for Medicare coverage, we also have $\alpha_{111} > \alpha_{101}$.

Then the triple-difference estimator can be written as:

$$\begin{aligned} DDD &= \alpha_{111} (Y_{111}^1 - Y_{111}^0) - \alpha_{101} (Y_{101}^1 - Y_{101}^0) \\ &\quad + (Y_{111}^0 - Y_{110}^0) - (Y_{011}^1 - Y_{010}^1) \\ &\quad - (Y_{101}^0 - Y_{100}^0) + (Y_{001}^1 - Y_{000}^1) \end{aligned}$$

The “parallel trends” assumption can be stated as the assumption that the second and third lines in the previous equality vanish and that $\alpha_{101} (Y_{101}^1 - Y_{101}^0) = 0$. While it is plausible that the second and third terms vanish, the $\alpha_{101} (Y_{101}^1 - Y_{101}^0)$, which reflects the effect of the expansions on the disabled without ESRD, is unlikely to vanish. Therefore DDD is biased by partial takeup of treatment ($\alpha_{111} < 1$) and the fact that some people without kidney disease are also treated ($\alpha_{101} > 0$). However, the bias can be signed if one assumes that the sign of the treatment effect is the same regardless of kidney disease status, in which case the triple-difference estimate will be biased towards zero unless the treatment effect of eligibility for people without kidney disease is significantly greater than the treatment for people with kidney disease.

In the difference-in-difference estimate that restricts to people with kidney disease, there is no bias from the fact that people without kidney disease are partially treated. One can write this estimator as:

$$DD_k = \alpha_{111} (Y_{111}^1 - Y_{111}^0) + (Y_{111}^0 - Y_{110}^0) - (Y_{011}^1 - Y_{010}^1)$$

Assuming that the parallel trends assumption holds, then DD_k provides a scaled estimate of the causal effect of Medicare eligibility for people with kidney disease. In the DD_k estimator, the parallel trends assumption implies that in the absence of the ESRD expansion, trends in mortality would have progressed along similar paths following the expansion for people over and under 65 years of age. While there are reasons to doubt this assumption, due to the fact that renal replacement therapy was generally more suited to younger, rather than older, people, it is unclear why there would be a sudden change at age 65, as would be required to bias my estimates.

It is tempting to also consider difference-in-difference models that compare people with and without kidney disease who are under 65, but such a model would be subject to the same biases as the triple difference model since some people without kidney disease also became eligible for Medicare coverage following the 1973 expansions.

3.2.2 Event Study and Difference-in-Difference Models

I first consider analyses that use age and time variation separately. To do so, I estimate event-study models of the form:

$$\begin{aligned}
 y_{iatgd} &= \beta_1^{65} Kidney_d + \beta_2^{65} Post_t + \beta_3^{65} Kidney_d \times Post_t \\
 &+ \sum_{a' \neq 65} (\beta_1^{a'} Kidney_d + \beta_2^{a'} Post_t + \beta_3^{a'} Kidney_d \times Post_t) \mathbf{1}_{[a=a']} \\
 &+ X_{ig} \Gamma_1 + \tau_t + \alpha_a + \varepsilon
 \end{aligned} \tag{1a}$$

and

$$\begin{aligned}
 y_{iatgd} &= \beta_1^{1971} Kidney_d + \beta_2^{1971} Under65_a + \beta_3^{1971} Kidney_d \times Under65_a \\
 &+ \sum_{t' \neq 1971} (\beta_1^{t'} Kidney_d + \beta_2^{t'} Under65_a + \beta_3^{t'} Kidney_d \times Under65_a) \mathbf{1}_{[t=t']} \\
 &+ X_{ig} \Gamma_1 + \tau_t + \alpha_a + \varepsilon
 \end{aligned} \tag{1b}$$

Where y_{iatgd} is the outcome—type of insurance coverage, amount of health care utilization, or deaths per 100,000 people—for person i (I only have person-level data on insurance coverage), who belongs to age group a in time period t , where time is measured in half-year increments, gender g , and cause of death d , $Kidney_d$ is a dummy for deaths due to kidney disease, $Post_t$ is a dummy for the ESRD period, which takes the value of 1 for time periods after July 1, 1973, $Under65_a$ is an indicator that a is less than 65, X_{ig} is a vector of controls including fixed effects for each demographic group, τ_t and σ_a are year and age fixed effects, respectively. The coefficients β_i^a and β_i^t are the age or period-specific coefficients on kidney disease, the post period (or being under 65), and their interaction.

I then summarize the results of these event studies using a triple-difference estimator, which is subject to bias from the SSDI program, and a difference-in-differences estimator that is unbiased, but may also be less precise. The triple-difference model can be written as:

$$\begin{aligned}
 y_{iatgd} &= \beta_1 Kidney_d + \beta_2 Post_t + \beta_3 Under65_a + \beta_4 Kidney_d \times Post_t \\
 &+ \beta_5 Kidney_d \times Under65_a + \beta_6 Post_t \times Under65_a \\
 &+ \beta_7 Kidney_d \times Post_t \times Under65_a + X_{ig} \Gamma_1 + \tau_t + \alpha_a + \varepsilon
 \end{aligned} \tag{2}$$

And the corresponding difference-in-differences estimator is:

$$y_{iatgd} = \alpha_1 Post_t + \alpha_2 Under65_a + \alpha_3 Post_t \times Under65_a + X_{ig} \Gamma_1 + \tau_t + \alpha_a + \varepsilon \tag{3}$$

The previous discussion of identification in this setting implies that $|\alpha_3| \geq |\beta_7|$, assuming that treatment effects in the ESRD expansion are comparable in size, or larger, than treatment effects of the SSDI expansion.

Standard errors for all models are clustered on age and time, unless otherwise specified, using `clus_nway.ado` (Cameron et al., 2011; Kleinbaum et al., 2013). I cluster on age to be consistent with the recommendations in Lee and Lemieux (2010) and Lee and Card (2008). I cluster on time based on recent results in Hausman and Rapson (2017).

4 Effect of the Medicare Expansion

4.1 Health Insurance

[Figure 1 about here.]

I first consider the effect of the Medicare expansion on health insurance coverage among people with kidney disease. Figure 1 presents an event study of the change in any insurance coverage (panel A) and Medicare coverage (panel B) using the triple-difference version of equation (1b).⁴ Prior to the Medicare expansion, the probability that an individual with kidney disease had any form of insurance coverage was increasing from 1968 to 1970, but stable from 1970 to 1972. However following the expansion there was no appreciable increase in insurance coverage, on average, for people with kidney disease. Medicare coverage, by contrast, increased significantly by 1978 with the bulk of the increase in Medicare coverage happening at older ages (panel D). Conversely, the ESRD expansion appears to have increased coverage somewhat for people close to the age 65 cutoff, but there was also a noticeable increase in insurance coverage for people around 40 years of age (panel C).

The results in figure 1 provides some support for the “parallel trends” assumption underlying differences-in-differences estimators. The fact that there was an increase in the point estimates from 1968 to 1970 for the probability of having any insurance is concerning, but this trend does not continue into 1972. I find no indication of a time trend in Medicare coverage. By age, panels C and D demonstrate that insurance coverage for people over 65 was not appreciably affected by the Medicare expansion.

[Table 2 about here.]

Consistent with the event study in figure 1, I find no evidence that the ESRD expansion increased insurance coverage among people with kidney disease (table 2, column 1). However, there was a 23 to 30 percentage point increase in coverage for people close to the age 65 cutoff (column 2). The increase in insurance coverage in models with age trends is slightly smaller than the increase in Medicare coverage (columns 3 and 4), which is consistent with either a degree of crowd-out or “doubling-up” of private and public insurance coverage. I find only modest evidence of a decrease in private insurance coverage associated with the ESRD expansion, but a

⁴The figure using the difference-in-difference is similar, but less precisely estimated.

large decrease in the share of people who reported only private insurance coverage. This final change—the reduction in reports of only private insurance—is indicative of people using both Medicare and private insurance coverage simultaneously. This kind of doubling up of insurance coverage provided additional benefits to people with ESRD since private insurance plans at the time typically did not cover dialysis or renal transplantation, hence adding Medicare coverage represented a significant improvement in insurance coverage for people with kidney disease.

Comparing the DDD and DD coefficients provides support for the fact that the Medicare expansion’s effect on people without kidney disease appears to bias my DDD estimates towards zero for having any insurance coverage and for Medicare coverage. This bias is what one would expect if the Medicare expansion also affected insurance coverage for some people without kidney disease, in this case by providing coverage to the long-term disabled without kidney disease.

The row labeled “Agg. Effect” presents the average aggregate effect of the Medicare expansion on insurance coverage using the triple difference or difference-in-difference coefficient, as appropriate. The average aggregate effect is the average of the annual total of the sampling weight for people between 45 and 64 years of age with kidney disease multiplied by the DDD or DD coefficient and indicates how many people with kidney disease gained or lost insurance coverage as a result of the Medicare expansion. These results indicate that as few as 5400 people or as many as 58000 people with kidney disease gained insurance coverage as a result of the expansion, although only the higher estimate is based on a statistically significant coefficient. Using data on Medicare coverage, I find a large increase in coverage of between 37000 and 66000 people having Medicare coverage. My estimates for Medicare coverage are large and are, in fact, larger than what Medicare trustees reported for the total number of people with ESRD, whether they became eligible solely due to having ESRD or because they were disabled. The fact that my implied increase in Medicare coverage is larger than the estimate from Medicare trustees should not be surprising since, in order to have sufficient data, I am applying a far more relaxed definition of kidney disease than is used by Medicare itself. The aggregate effect estimated using the DD coefficient is consistently larger than the estimate from the DDD estimate, which is what I had hypothesized based on the fact that some people without kidney disease were also gaining access to Medicare coverage.

As a specification check, online appendix table B1 presents results from “donut” regressions that exclude people within five years of turning 65. These donut estimates are, in general, consistent with my main specifications, particularly for models without age trends.

4.2 Health Care Utilization

[Figure 2 about here.]

Figure 2 presents triple difference estimates for the number of physician visits. The estimates in panel A are extremely noisy, both in terms of the standard error, but also in the point estimate itself, with relatively large amount of variation in the point estimate both before and after the expansion took effect. However, visually it appears that there may have been an increase in physician visits after, versus before, the Medicare expansion for people with kidney disease. Panel B demonstrates that any increase in physician visits affected virtually all ages below 65 years of age and there was essentially no effect on physician visits for people over 65 years of age.

Consistent with the event studies, column 9 of table 2 demonstrates that the ESRD expansion was associated with a 18 to 25 percent increase in physician visits in models that do not control for age trends. Models that do control for age trends (column 10) yield larger estimates, which is consistent with the age profile of the change in physician visits from panel B of figure 1. In aggregate, my DDD estimates indicate that there were an additional 400000 to 600000 physician visits per year among people with kidney disease, or an additional 2.3 to 3.5 visits per person with kidney disease (“Avg. Effect” row).⁵

As was the case with my insurance estimates, my difference-in-difference results are generally comparable, although larger, than my triple-difference estimates and this extends to the aggregate and average effects of the expansion on doctor visits.

These results are essentially unchanged in donut regressions (online appendix table B1, columns 9 and 10).

4.3 Mortality Effects

4.3.1 Comparisons within the United States

[Figure 3 about here.]

Figure 3 plots event studies for kidney disease mortality using the underlying cause of death, where the event studies are based on triple-difference estimates. Panel A indicates that there was a reduction in mortality due to kidney disease in 1973 and visually, this reduction was larger than the potential downward trend in kidney disease mortality prior to 1973. Panel B demonstrates that there was a relative increase in kidney disease mortality among people 65 years of age and older following the ESRD expansion and indicates that there was a strong age trend in the mortality change following the ESRD expansion, which justifies focusing on specifications that include age trends.

[Table 3 about here.]

In triple difference models based on equation (2) and difference-in-differences estimates based on equation (3) I find that the ESRD program reduced mortality

⁵There were approximately 180,000 people who met my definition of kidney disease per year between 45 and 64 years of age.

from kidney disease by 36.3 to 37.3 log points in a model that does not include age trends. This estimate is, at first blush, implausibly large and reflects the age trends seen in panel B of figure 3. Including age trends (column 2) yields smaller estimates of a 7.3 to 7.9 log point reduction in mortality. In models with age trends, I also find that the DD estimate is larger in magnitude (although not significantly so) than the DDD estimate, which is what one would expect from the lower level of Medicare eligibility among people without kidney disease.

Because there are many potential diagnoses that may indicate a death due to kidney disease, in columns (5) and (6) I present results using the “broad” definition of kidney disease. The broad results are qualitatively similar and also indicate a reduction in mortality based on kidney disease as the underlying cause of death, but not when kidney disease is defined using both underlying and contributing cause of death codes.

Lastly, the ESRD program, in particular, was targeted at people with chronic kidney disease, so in columns (7) and (8) I restrict my definition of kidney disease to people who died of chronic kidney disease, based on the codings in appendix table A. The chronic estimates indicate that the Medicare expansion was associated with a reduction in deaths due to chronic kidney disease, although three of the estimates are only significant at the ten percent level.

Table 3 also presents the change in the mortality rate and the implied number of deaths averted, based on population data from 1973 and the average mortality rate due to kidney disease before the Medicare expansion. Under the narrow definition and without age trends, the Medicare expansion appears to have reduced the mortality rate by 1.9 to 2.3 deaths per 100,000 for a total of between 700 and 900 fewer deaths per year due kidney disease. However, these estimates are significantly narrowed in models that include age trends (columns 2 and 4) to a reduction of 0.5 to 1.0 deaths per 100,000 or 180 to 390 fewer deaths. The fact that there is a greater reduction in the number of deaths using both underlying and contributing causes of death, versus just underlying causes of death, indicates that at least part of the reduction in mortality that I observed in column (2) is not due to a change in coding practices in which kidney disease was less likely to be listed as an underlying cause of death, but more likely to be listed as a contributing cause of death.

Repeating the change in mortality analysis for both the broad and chronic definitions yields two interesting results. First, the vast majority of the reduction in kidney disease deaths is arising from fewer deaths due to chronic disease, particularly when using both underlying and contributing causes. Second, while there is a larger reduction in mortality under the broad definition when I only look at underlying cause of death codes, I actually find that there were fewer deaths averted under the broad definition than the narrow definition when I use both underlying and contributing causes of death.

The fact that there are fewer deaths averted using underlying and contributing cause of death codes versus just the underlying cause of death codes under the broad

definition is unexpected. One might expect that under a more relaxed definition of kidney disease mortality the opposite would occur. The break in the pattern between underlying and underlying and contributing mortality does not reflect a change in coding practices since there were no relevant changes in the coding manuals published by the National Center for Health Statistics during this time period. However, what appears to be happening is that there is essentially no change in the number of deaths that list one of the more tenuously related cause of death codes as a contributing cause of death. Given the set of broad cause of death codes, it is not surprising that there is a smaller change in the number of deaths that list, for example, hypertension as a contributing cause of death.

The mortality reductions in panel A of table 3 can be combined with the estimated change in insurance coverage from table 2 to infer how large an effect insurance coverage may have on kidney disease mortality. Using either the increase in insurance coverage from column (2) or the increase in Medicare coverage from column (4) implies that there was a reduction of 0.4 percentage points (any insurance) or 0.33 to 0.35 percentage points (using the increase in Medicare coverage) in the probability of dying from kidney disease associated with insurance coverage. These estimates are about three times larger than the local average treatment effect of Medicaid coverage estimated by the Oregon Health Insurance Study (column (3) of table IX in Finkelstein et al., 2012).

Alternatively, one can compute the elasticity of mortality with respect to insurance coverage. From table 2, the percentage change in insurance coverage is 0.31 based on the DDD estimate for any insurance coverage and 0.41 using the DD estimate. Using the point estimates in table 3 yields an elasticity of mortality using the narrow, underlying definition of -0.23 using the DDD and -0.19 using the DD estimate, which is twice as large as the elasticity from the Oregon Health insurance Study.⁶

There are two main threats to the validity of my results that are unique to mortality data. First, there is a reverse “harvesting” effect, in which people who would have died of kidney disease in the absence of the program are able to survive until they turn 65 after the program. The implication of this kind of harvesting is that the mortality rate among people 65 and older will be overstated. I am able to test for this possibility by re-running my underlying models while excluding people between 60 and 70 years of age (panel B). In these donut regressions, my results are essentially unchanged and, in fact, my estimated mortality reductions become larger. This is inconsistent with reverse harvesting, which would predict that the mortality reductions would be smaller in magnitude when I exclude people between 60 and 70 years of age.

⁶To calculate the elasticity, I first calculated the fraction of the controls who died by the end of the study (0.8 percent) to get an implied percentage change in mortality of 16.25% ($\frac{0.0013}{0.008}$). The first-stage statistics (table III) implies a percentage change in Medicaid coverage of 182%. The resulting elasticity is 0.09 ($\frac{0.1625}{1.816}$).

The second threat is that people who do not die of kidney disease will die of something else. This “competing risk” effect is well known in economics and epidemiology and cannot be resolved without imposing assumptions on the processes that determine mortality (Honoré and Lleras-Muney, 2006). The bias due to competing risks is similar to the bias from harvesting, but now it is the mortality rate due to non-kidney causes that is inflated. Notably, competing risks can only bias my estimates if there is, in fact, an effect of the Medicare expansion on kidney disease mortality. In the absence of such a reduction, there is no reason to expect to find a competing risk bias. I can address the bias from competing risks by restricting my data to deaths due to kidney disease, in other words the DD results are not subject to competing risks. My DD results demonstrate that any bias from competing risks is small since my DD estimates are, in general, larger in magnitude than the DDD estimates (which is also the relationship one would expect to hold if the treatment effect of Medicare eligibility was of the same sign for people with and without kidney disease).

The online appendix presents results from a log-linear OLS specification, which are qualitatively similar (table B2). The online appendix also provides robustness tests of the triple and double-difference results by varying the range of ages included (online appendix figure A1) and varying the age and time controls that are included (table B3).

In the online appendix (table B4), I also consider the potential confounding effect of the introduction of the Supplemental Security Income (SSI) program in 1974. The SSI program provides cash transfers to low-income people who are aged, blind, or disabled and, in most states, also provides access to Medicaid coverage. To test if the SSI program is confounding my estimate of the effect of the SSDI program, I interacted the triple difference coefficients with (demeaned) shares of people in an age-gender-state cell who reported either Social Security or SSI income in the March CPS from 1977 to 1979 (covering years 1976 to 1978). In a separate specification, I interacted the triple difference coefficients with indicators for two factors that states can use to discourage enrollment in Medicaid—using more stringent eligibility criteria and requiring a separate Medicaid application. I find no evidence that any of these interactions are statistically significant, indicating that the SSI program is not driving the differential mortality reduction for kidney disease, relative to other causes of death.

4.3.2 Comparisons with Other OECD Countries

[Figure 4 about here.]

Figure 4 plots event-study estimates of the change in kidney disease mortality in the United States, relative to other OECD countries by either year (panel A) or age (panel B). Over time, there is a pronounced reduction in kidney disease mortality for people under 65 in the United States in 1973 that was not observed in other

countries. However, there is also some evidence of a trend in kidney disease mortality in the United States towards fewer people under 65 dying from kidney disease, although with one exception, all of the confidence intervals before 1972 include 0. Despite the possible violation of the parallel trends assumption, there is still evidence of a substantial reduction in kidney disease mortality beginning in 1973. Results by age (panel B) are also suggestive of a reduction in kidney disease mortality, although there appears to be a reduction in mortality among 65-70 year olds, relative to people 70 and older, in the data as well.

[Table 4 about here.]

Going from the event-study estimates in figure 4 to triple and quadruple difference results, I find that the ESRD program was associated with a four to eight log point reduction in mortality from kidney disease, depending upon the specification and sample (table 4). This reduction in mortality is robust to including country fixed effects, interacting country fixed effects with either kidney disease or an indicator for 1974 or later (the post dummy takes the value 0.5 in 1973), and including year-by-kidney disease indicators, which accounts for innovations in the treatment of kidney disease. These results are also similar in magnitude to my results using the narrow definition of kidney disease and underlying cause of death codes in the US mortality data, which is the most comparable specification.

5 Mechanisms

The ESRD expansion may have affected health through two classes of mechanisms. First, by lowering the cost of accessing treatment, health insurance may have increased demand for renal replacement services (dialysis and kidney transplantation), which would have been otherwise unaffordable. This mechanism implies that there may be an “access motive” to purchase health insurance in the sense of Nyman (2003, 1999a,b) and, in essence, reflects the fact that the Medicare expansion provided a large in-kind transfer from healthy people to those with ESRD.

The second class of mechanisms relate to changes in the supply of renal replacement services. The expansion did not merely shift the demand curve outward, but it also guaranteed payment for treatment services, which reduced the risk of investing in renal replacement services. In much the same way that the original introduction of Medicare stimulated entry by hospitals and increased technology adoption (Finkelstein, 2007), the ESRD expansion may have increased adoption and entry of renal replacement services across the country.

5.1 Access to Care

[Table 5 about here.]

In order to test if access to care was an important contributor to the reduction in mortality associated with the ESRD expansion, I augmented equation (2) with interactions between the triple-difference variables and measures of the number of dialysis clinics and transplant programs per capita. Table 5 presents the results of this analysis. Panel A demonstrates that living in a state with more dialysis clinics in 1971 was associated with a significantly larger decline in kidney disease mortality, but there was no effect of living in a state with a transplant program. These results persist, even after I include indicators for the presence of Veteran’s Administration dialysis clinics and transplant programs (panel B).

The lack of evidence that transplant programs affect local mortality is not surprising since transplant programs require fewer visits than dialysis clinics. Therefore patients may be willing to travel long distances in order to get a kidney transplant, meaning that the number of programs in a state is not the most relevant metric affecting their survival.

The implication of these results is that either the ESRD program increased the number of dialysis clinics in states that already had a large number of clinics, relative to population, or that the program facilitated access to the existing clinic network. In the next subsection, I test if the number number of dialysis clinics per capita after the expansion increased more in areas with more dialysis clinics per capita in 1971, or if there was greater entry in areas with fewer dialysis clinics per capita.

5.2 Entry of Treatment Facilities

The ideal data with which to test the entry hypothesis would involve regressing the change in treatment facilities on the number of people for whom dialysis or kidney transplantation was appropriate. However, such data are not available. Instead, I use the mortality rate due to kidney disease as a proxy. The idea behind this proxy is that in areas with a higher mortality rate there are likely to be more people for whom treatment is appropriate. Therefore, to test the entry hypothesis, I estimate the following model:

$$\ln E \left[\frac{y_s^{1975}}{pop_s^{1975}} \right] = \alpha_0 + \alpha_1 \ln \left(\frac{y_s^{1971}}{pop_s^{1971}} \right) + \alpha_2 \ln Mort_{s,<65}^{1971} + \alpha_3 \ln Mort_{s,\geq 65}^{1971} + \alpha_4 \mathbf{1} [y_s^{1971} = 0] \quad (4)$$

Where the model is estimated as a Poisson regression, s denotes the state, superscripts refer to the year to which the data refer, y_s^t is either the number of dialysis clinics or transplant programs in state s at time t , $Mort_{s,g}^t$ is the kidney disease mortality rate using the “narrow” definition with deaths to attributed kidney disease based on the underlying cause of death codes in state s , year t , for age group g (either under 65 or 65 and older), and pop_s^t is the population in state s and year t . α_1 tests if the measure of treatment programs in a state is converging across the country depending on whether or not the elasticity of 1975 treatment capacity with

respect to 1971 treatment capacity is greater than, less than, or equal to one. α_2 and α_3 test if treatment capacity is responsive to the burden of disease in the area since areas with a greater burden of disease will have a higher mortality rate due to kidney disease. *A priori* one would expect $\alpha_2 > 0$ and $\alpha_3 \cong 0$ as indicators that the Medicare expansion, since it affected people under 65 years of age, encouraged entry.

[Table 6 about here.]

Table 6 demonstrates that the number of dialysis clinics (columns 1-3) was converging over time since the coefficient on 1971 treatment capacity is less than one. In other words, states with comparatively few dialysis clinics, relative to population, in 1971 experienced a more rapid rate of increase than did states with more dialysis clinics per capita in 1971. Furthermore, there is evidence that mortality among people under 65 served to increase the number of clinics in a state in 1975, which is consistent with the Medicare expansion encouraging entry of new dialysis clinics. Column 4-6 demonstrate that this pattern was weaker for transplant programs, with the elasticity of transplant programs per capita being significantly closer to 1 than was the case for dialysis clinics. There is also no evidence that entry of transplant programs was correlated with the 1971 burden of disease among people under 65 years of age, but rather kidney disease mortality among people 65 and over was more strongly correlated with the number of transplant programs in a state in 1975.

The fact that the Medicare expansion promoted greater entry of dialysis clinics than transplant programs is consistent with differences in how these two forms of treatment are used. Dialysis clinics require that patients return frequently for treatment since the typical treatment regimen may include as many as five treatments per week, as a result proximity to a dialysis clinic is important, hence one would expect to see a large increase in dialysis clinics. On the other hand, kidney transplant programs require fewer visits so that patients may be willing to travel long distances in order to get a kidney transplant, meaning that there is a weaker incentive for new transplant programs to enter in response to patient demand.

6 Welfare Implications

These results provide some insight into the welfare consequences of the Medicare expansion among people with kidney disease. The first margin to study are the implications of my results for the productivity of moral-hazard induced care. One typically thinks of moral-hazard induced care as inefficient since it is care that the consumer was unwilling to pay for at the offered price (Pauly, 1968). However, one can recast this framework in terms of the marginal health product of health care and a consumer's willingness to pay for a unit of health. In this framework, a consumer's willingness to pay for health care is decreasing because either the marginal health product is decreasing or her valuation of a unit of health is decreasing. Assuming

that a person’s value of a unit of health is fixed (or at least unlikely to change significantly) then the downward slope of demand curves for health care (and the resulting welfare losses from moral hazard) come from the decreasing marginal product of health care.

In this paper, I provide suggestive evidence of an increase in utilization of dialysis facilities. First, I find an increase in self-reported physician visits in the NHIS, which includes dialysis care. Second, the reduction in kidney disease mortality was larger in areas with more dialysis clinics in 1971. Collectively, these results suggest that the increase in dialysis clinic visits had a positive marginal health product. Whether or not this health impact was large enough to eliminate the welfare cost of the increase in consumption depends on the size of the health improvement.

I can quantify the size of the health improvement by computing the change in survival associated with the program and, from there, calculating the number of life years saved due to the program. In order to estimate the survival gains, I began by computing the age- and gender-specific average mortality hazard due to kidney and non-kidney causes in the pre period. In order to compute counterfactual survival, I then multiplied the mortality hazards due to kidney disease for people under 65 years of age by the exponentiated triple difference or difference-in-difference coefficient from the models in table 3. In order to combine these mortality hazards into a single hazard that I could use to calculate survival, I assumed that latent survival durations from kidney and non-kidney causes were independent so that the mortality hazard at age a is the sum of the cause specific mortality hazards at age a .⁷ I computed survival from age 45 as the sum of the cumulative survival probabilities⁸ and computed the differences from the survival durations implied by the pre-period mortality rates. I then converted these differences, which are representative of the effects on a 45 year old, into population-level estimates by multiplying by the population of white 45 years olds in 1973, which yields an estimate of the number of life years saved by the Medicare expansion’s effect on kidney disease mortality and, therefore, the productivity of the induced health care utilization.

[Table 7 about here.]

Table 7 presents the results of this analysis. In column (2), which reports results using the narrow definition and with age trends, in the first panel I report that life expectancy from age 45 is 18.36 years up to age 65 and almost 29 years to age 85 (I am unable to calculate subsequent mortality hazards since I do not have denominator data for people 85 and older). Using the triple-difference coefficients there is almost no change in survival—life expectancy rose by 0.001 years to age 64 and by 0.002

⁷Assuming that survival durations are independent is a strong assumption, but some assumption is needed in order to evaluate treatment effects in this kind of a competing risks framework (Honore and Lleras-Muney, 2006).

⁸Suppressing all subscripts other than age, if the mortality rate at age a is denoted by m_a then the survival duration is $S = \sum_{a=45}^{84} [\exp(\sum_{a'=45}^a \log(1 - m_{a'}))]$.

years to age 84. However, these estimates are for the entire population while only a small minority actually has kidney disease. When I scale these estimates by the number of white and non-white 45 year olds, I find that the expansion saved between 2200 and 5100 life years, depending on the age cutoff used. I find larger savings using the difference-in-difference estimate to construct the counterfactual mortality rates. Applying a value of \$100,000 to a life year, the results in column (2) imply that the mortality benefits of the Medicare expansion due to changes in kidney disease mortality were worth between \$220 million and \$670 million per year. Spending on this population in a single year was around \$750 million indicating that the program cannot be justified solely based on its effects on kidney-related mortality.⁹ However, using some of my more relaxed specifications (e.g. including contributing causes of death) implies that the value of the life years saved may exceed \$1 billion, indicating that it is possible that this expansion yielded benefits in excess of costs, assuming that each life year was worth \$100,000 and that other costs associated with the program (e.g. spending on other services) are not too large.

Across most of the remaining specifications, I find evidence of an increase in survival, with estimates using underlying causes of death and age trends indicating that the expansion saved between 2500 and 14000 life years; using contributing cause of death codes as well yields a somewhat broader range, though the increase is not as dramatic as with the narrow definition of kidney disease.

7 Conclusions

In this paper, I estimated the causal effect of the 1973 Medicare expansions affected people with kidney disease. In aggregate the expansion increased insurance coverage and physician visits for people with kidney disease. I also document a significant reduction in mortality due to kidney disease that was robust to a variety of specification checks and alternative definitions of kidney disease.

I identify two mechanisms for my results. The first mechanism is that the increase in insurance coverage provided access to treatment that was otherwise unavailable (Nyman, 2003, 1999a,b). Consistent with this mechanism, I find larger reductions in kidney disease mortality for people under 65 in areas that had more dialysis facilities in 1971. An important implication of this mechanism is that there is a large liquidity effect in the demand for medical care, in which case the welfare loss from moral hazard may be significantly reduced.

I also find evidence in support of a second, supply-side, mechanism by which the Medicare expansion lead to increased entry of dialysis clinics. Specifically, I find that having a higher mortality rate due to kidney disease among people under 65 in 1971 is correlated with having more dialysis clinics per capita in 1975. I do not

⁹I assumed that there were 11500 beneficiaries between 45 and 65 years of age based on the age distribution of people who were eligible for Medicare only because of ESRD (people who did not also have a long-term disability) and that spending per person was \$64,500 per year in 2015 dollars.

find a comparable effect for transplant programs, which is consistent with transplant programs competing across larger geographic areas, while dialysis clinics compete in more local markets.

My results contribute to a large literature on the effects of public insurance programs (e.g. Currie and Gruber, 1996a,a; Cutler and Gruber, 1996; Finkelstein, 2007; Finkelstein and McKnight, 2008; Finkelstein et al., 2012; Goodman-Bacon, 2017; Gruber and Simon, 2008). However, a distinctive feature of my results, relative to others in the literature, is that the program that I study conditions coverage on being in poor health. As a result, the benchmark for evaluating this program is somewhat different than for other insurance expansions since an effect on mortality that may seem large among a population that was not selected on the basis of ill health, may be much more plausible in the context of a program that explicitly conditioned eligibility on people having an expected survival of days or weeks following diagnosis with ESRD.

References

- Shana Alexander. They decide who lives, who dies. *Life Magazine*, 53(19):102–125, November 1962. URL <https://repository.library.georgetown.edu/handle/10822/762327>.
- Robert M. Ball. Social Security Amendments of 1972: Summary and Legislative History. *Social Security Bulletin*, 36(3):3–25, March 1973.
- Silvia Helena Barcellos and Mireille Jacobson. The Effects of Medicare on Medical Expenditure Risk and Financial Strain [†]. *American Economic Journal: Economic Policy*, 7(4):41–70, November 2015. ISSN 1945-7731, 1945-774X. doi: 10.1257/pol.20140262. URL <http://pubs.aeaweb.org/doi/10.1257/pol.20140262>.
- A. Colin Cameron, Jonah B. Gelbach, and Douglas L. Miller. Robust Inference With Multiway Clustering. *Journal of Business and Economic Statistics*, 29:238–249, April 2011. ISSN 0735-0015, 1537-2707. doi: 10.1198/jbes.2010.07136. URL <http://pubs.amstat.org/doi/abs/10.1198/jbes.2010.07136>.
- David Card, Carlos Dobkin, and Nicole Maestas. The Impact of Nearly Universal Insurance Coverage on Health Care Utilization: Evidence from Medicare. *American Economic Review*, 98(5):2242–2258, December 2008. ISSN 0002-8282. doi: 10.1257/aer.98.5.2242. URL <http://www.aeaweb.org/articles.php?doi=10.1257/aer.98.5.2242>.
- David Card, Carlos Dobkin, and Nicole Maestas. Does Medicare Save Lives? *Quarterly Journal of Economics*, 124(2):597–636, May 2009. doi: 10.1162/qjec.2009.124.2.597. URL <http://www.mitpressjournals.org/doi/abs/10.1162/qjec.2009.124.2.597>.

- Kenneth Y. Chay, Daeho Kim, and Shailender Swaminathan. Medicare's Impact on Hospital Insurance, Hospital Utilization, and Life Expectancy: The First 25 Years. 2017.
- Congressional Research Service. Hemodialysis and Kidney Transplantation: Practice and Policy in Total Organ Failure. Technical report, Congressional Research Service, Washington, D.C., November 1971.
- Janet Currie and Jonathan Gruber. Health Insurance Eligibility, Utilization of Medical Care, and Child Health. *The Quarterly Journal of Economics*, 111(2): 431–466, May 1996a. ISSN 00335533. doi: 10.2307/2946684. URL <http://www.jstor.org/stable/2946684>.
- Janet Currie and Jonathan Gruber. Saving Babies: The Efficacy and Cost of Recent Changes in the Medicaid Eligibility of Pregnant Women. *The Journal of Political Economy*, 104(6):1263–1296, December 1996b. ISSN 00223808. doi: 10.2307/2138939. URL <http://www.jstor.org/stable/2138939>.
- David M Cutler and Jonathan Gruber. Does Public Insurance Crowd Out Private Insurance. *The Quarterly Journal of Economics*, 111(2):391–430, May 1996. ISSN 00335533. URL <http://links.jstor.org/sici?sici=0033-5533%28199605%29111%3A2%3C391%3ADPICOP%3E2.0.CO%3B2-S>.
- Gary V. Engelhardt and Jonathan Gruber. Medicare Part D and the Financial Protection of the Elderly. *American Economic Journal: Economic Policy*, 3(4):77–102, November 2011. ISSN 1945-7731. doi: 10.1257/pol.3.4.77. URL <https://www.aeaweb.org/articles?id=10.1257/pol.3.4.77>.
- Amy Finkelstein. The Aggregate Effects of Health Insurance: Evidence from the Introduction of Medicare*. *Quarterly Journal of Economics*, 122(1):1–37, February 2007. doi: 10.1162/qjec.122.1.1. URL <http://dx.doi.org/10.1162/qjec.122.1.1>.
- Amy Finkelstein and Robin McKnight. What did Medicare do? The initial impact of Medicare on mortality and out of pocket medical spending. *Journal of Public Economics*, 92(7):1644–1668, July 2008. ISSN 0047-2727. doi: 10.1016/j.jpubeco.2007.10.005. URL <http://www.sciencedirect.com/science/article/B6V76-4R3C010-1/2/f7d778844604aa7092b85fc128f078a1>.
- Amy Finkelstein, Sarah Taubman, Bill Wright, Mira Bernstein, Jonathan Gruber, Joseph P. Newhouse, Heidi Allen, and Katherine Baicker. The Oregon Health Insurance Experiment: Evidence from the First Year*. *The Quarterly Journal of Economics*, 127(3):1057–1106, August 2012. ISSN 0033-5533, 1531-4650. doi: 10.1093/qje/qjs020. URL <http://qje.oxfordjournals.org/content/127/3/1057>.

- Andrew Goodman-Bacon. Public Insurance and Mortality: Evidence from Medicaid Implementation. *Journal of Political Economy*, 126(1):216–262, October 2017. ISSN 0022-3808. doi: 10.1086/695528. URL <https://www.journals.uchicago.edu/doi/abs/10.1086/695528>.
- Jonathan Gruber and Kosali Simon. Crowd-out 10 years later: Have recent public insurance expansions crowded out private health insurance? *Journal of Health Economics*, 27(2):201–217, March 2008. ISSN 0167-6296. doi: 10.1016/j.jhealeco.2007.11.004. URL <http://www.sciencedirect.com/science/article/B6V8K-4R7NPWM-2/2/56ca97e166ebf606c652a9fcf6ce5d7d>.
- Catherine Hausman and David S. Rapson. Regression Discontinuity in Time: Considerations for Empirical Applications. Working Paper 23602, National Bureau of Economic Research, July 2017. URL <http://www.nber.org/papers/w23602>. DOI: 10.3386/w23602.
- Bo E. Honoré and Adriana Lleras-Muney. Bounds in Competing Risks Models and the War on Cancer. *Econometrica*, 74(6):1675–1698, November 2006. ISSN 1468-0262. doi: 10.1111/j.1468-0262.2006.00722.x. URL <http://onlinelibrary.wiley.com/doi/10.1111/j.1468-0262.2006.00722.x/abstract>.
- Kidney Disease Program. *Kidney disease services, facilities, and programs in the United States*. Regional Medical Programs Services, Kidney Disease Control Program; National Kidney Foundation, Rockville, Md., New York, 1971. URL <https://catalog.hathitrust.org/Record/011324423>.
- Adam M. Kleinbaum, Toby E. Stuart, and Michael L. Tushman. Discretion Within Constraint: Homophily and Structure in a Formal Organization. *Organization Science*, 24(5):1316–1336, February 2013. ISSN 1047-7039. doi: 10.1287/orsc.1120.0804. URL <http://pubsonline.informs.org/doi/abs/10.1287/orsc.1120.0804>.
- David S. Lee and David Card. Regression discontinuity inference with specification error. *Journal of Econometrics*, 142(2):655–674, February 2008. ISSN 0304-4076. doi: 10.1016/j.jeconom.2007.05.003. URL <http://www.sciencedirect.com.ezp-prod1.hul.harvard.edu/science/article/B6VC0-4NT9GJ9-2/2/07619f69a2aa922fafd1cfaf2dbd36c6>.
- David S Lee and Thomas Lemieux. Regression Discontinuity Designs in Economics. *Journal of Economic Literature*, 48(2):281–355, June 2010. ISSN 0022-0515. doi: 10.1257/jel.48.2.281. URL <http://www.aeaweb.org.ezp-prod1.hul.harvard.edu/articles.php?doi=10.1257/jel.48.2.281>.
- Myoung-jae Lee and Changhui Kang. Identification for difference in differences with cross-section and panel data. *Economics Letters*, 92(2):270–276, August


2006. ISSN 0165-1765. doi: 10.1016/j.econlet.2006.03.007. URL <http://www.sciencedirect.com/science/article/pii/S0165176506000802>.
- National Kidney Foundation. Your Kidneys: Master Chemists of the Body. Technical report, National Kidney Foundation, 2009. URL <https://www.kidney.org/sites/default/files/docs/masterchemists.pdf>.
- John Nyman. *The Theory of Demand for Health Insurance*. Stanford Economics and Finance, 1 edition edition, 2003. ISBN 0-8047-4488-2.
- John A. Nyman. The economics of moral hazard revisited. *Journal of Health Economics*, 18(6):811–824, December 1999a. ISSN 0167-6296. doi: 10.1016/S0167-6296(99)00015-6. URL <http://www.sciencedirect.com/science/article/B6V8K-3Y9V2DT-7/2/b1ed50b860aa122df3093ee223e602bd>.
- John A. Nyman. The value of health insurance: the access motive. *Journal of Health Economics*, 18(2):141–152, April 1999b. doi: 10.1016/S0167-6296(98)00049-6. URL <http://www.sciencedirect.com/science/article/B6V8K-3VKSDTM-1/2/c70c5a72fb864daa8a2a437e47ff1b74>.
- Mark V. Pauly. The Economics of Moral Hazard: Comment. *The American Economic Review*, 58(3):531–537, June 1968. ISSN 00028282. URL <http://www.jstor.org/stable/1813785>.
- Richard A. Rettig. The policy debate on patient care financing for victims of end-stage renal disease. *Law and Contemporary Problems*, 40(4):196–230, 1976. ISSN 0023-9186.
- Richard A. Rettig. Origin of the Medicare Kidney Disease Entitlement: The Social Security Amendments of 1972. In *Biomedical Politics*, pages 176–214. National Academies Press, Washington, D.C., January 1991. ISBN 978-0-309-04486-8. URL <http://www.nap.edu/catalog/1793>.
- Richard A. Rettig. Special Treatment — The Story of Medicare’s ESRD Entitlement. *New England Journal of Medicine*, 364(7):596–598, February 2011. ISSN 0028-4793. doi: 10.1056/NEJMp1014193. URL <http://dx.doi.org/10.1056/NEJMp1014193>.
- United States Department of Health and Human Services. National Center for Health Statistics. Multiple Cause of Death, 1968-1973, 2007a. URL <http://doi.org/10.3886/ICPSR03905.v2>.
- United States Department of Health and Human Services. National Center for Health Statistics. Multiple Cause of Death, 1974-1978, 2007b. URL <http://doi.org/10.3886/ICPSR03906.v2>.

Figures

Figure 1: Changes in Insurance Coverage from the Medicare Expansion


A: Any Insurance

B: Medicare



C: Any Insurance

D: Medicare




Source—National Health Interview Survey, even number years from 1970-1980

Notes—Sample restricted to people between 45 and 84 years of age. Points are year-by-under 65 years of age coefficients from a linear regression of an indicator for either any insurance or Medicare on year fixed effects (omitted 1970), an under 65 indicator, kidney disease status (using the “broad” definition), and their interaction. Confidence intervals are clustered on age.

Figure 2: Event Study Estimates of Changes in Health Care Utilization

A: # Doctor Visits—Year×Under 65×Kidney B: # Doctor Visits—Age×Post×Kidney




Source—National Health Interview Survey, 1969-1980.

Notes—Sample in panel A restricted to people between 45 and 84 years of age, except for people 65 years of age; panel B excludes observations from the second half of 1973 and the first half of 1974. Points in panel A are year-by-under 65 years of age-by-kidney disease coefficients from a regression of the number of doctor visits on year fixed effects (omitted 1970), an under 65 indicator, an indicator for kidney disease, and all two- and three-way interactions. Panel B presents point estimates for years of age interacted with a post dummy (after 1973) and kidney disease from a regressions of the number of doctor visits on age fixed effects, post, kidney disease, and all two- and three-way interactions; smoothed line is local polynomial estimate where estimates are weighted by the inverse of their standard errors. Estimates are from Poisson regressions. Confidence intervals in panel A based on covariance matrix that is clustered on age, while panel B uses heteroskedasticity robust standard errors. Kidney disease defined using the “broad” definition (see appendix table A).

Figure 3: Event Study Estimates of the ESRD Program and Mortality

A: Underlying Cause of Death by Year (1972 reference)

B: Underlying Cause of Death by Age (65 reference)




Source—Author’s analysis of Multiple Cause of Death Files, 1968-1998.


Notes—Points are coefficients on time periods from a Poisson regressions of the number of deaths per 100,000 in an age-gender-time period cell on time fixed effects (panels A and B) or age fixed effects (panels C and D) interacted with a dummy for kidney disease and either a dummy for being under 65 or after the second half of 1973. Kidney disease deaths classified using the underlying cause of death only in panels A and C and, in all panels, the “narrow” definition of kidney disease (see appendix table A). 95% confidence intervals are clustered on age and time.

Figure 4: Cross-Country Event Study Estimates of the ESRD Program and Mortality

A: Mortality Differences by Year



B: Mortality Differences by 5-Year Age Group



Source—Author’s analysis of the World Health Organization Mortality Database for 1968 through 1978.

Notes—Points in panel A are quadruple-difference estimates by year comparing mortality from kidney disease to all other deaths in the United States versus all other OECD member states that joined prior to 1969, relative to 1971 for the indicated age groups. Panel B reports quadruple-difference estimates by age group comparing before, versus after, the Medicare expansion, with versus without kidney disease, in the United States versus all other OECD member states. Standard errors are clustered on country.

Tables

Table 1: Enrollment, Spending, and Utilization in the ESRD Program

Year	Enrollment		Kidney Deaths		Spending		Utilization	
	Total	Under 65	Under 65	65 and Over	Total	Per enrollee	Transplants	Dialysis
1971			5335	7534				
1974	15993		4633	8949	1050.3	65673		
1975	22674	12702 ^a	4540	9491	1545.6	68164		
1976	28941	14721 ^a	4532	10597	2086.9	72110		
1977	35889	16514 ^a	4345	11008	2449.2	68243		
1978	43482	34828	4498	11973	2804.6	64500		
1979	52636	43031	3761	11966	3126.4	59397	4189	45565
1981	61930	47520	3761	13703	3723.7	60127	4898	58924
1986	93197	59570	3914	17851	6786.7	63646	8948	90886
1991	142510	83443	3395	17963	9704.2	56844	10037	144175
1996	255578		3433	20869	14141.8	55333	12219	215557

Source—Greenbook (various years), Annual Statistical Supplement to the Social Security Bulletin (various years), Multiple Cause of Death files, 1971-1996.

^a Enrollees eligible solely due to ESRD.

Notes—Enrollment based on enrollment in Medicare Part A, expenditures are for Medicare Parts A and B. Spending data have been inflated to 2015 using the CPI for urban workers. Utilization data are the number of transplants and number of enrollees dialyzed, respectively. Kidney deaths are based on chronic coding only, see appendix table A; the coding of kidney deaths changed between 1978 and 1979.

Table 2: Effect of the ESRD Program on Health Insurance and Health Care Utilization

	Any Insurance		Medicare		Any Private		Only Private		Doctor Visits	
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
DDD	0.021 (0.068)	0.226* (0.082)	0.144* (0.052)	0.260+ (0.126)	-0.084 (0.081)	0.056 (0.108)	-0.243* (0.081)	-0.205* (0.071)	0.181** (0.053)	0.266+ (0.159)
Agg. Effect ^a	5409	57732	36866	66329	-19614	12893	-62070	-52429	411298	632898
Avg. Effect ^b									2.28	3.50
N	147669	147669	118375	118375	188071	188071	118375	118375	371181	371181
DD	0.073 (0.069)	0.296** (0.092)	0.193** (0.056)	0.339* (0.131)	-0.164* (0.076)	-0.006 (0.123)	-0.238** (0.069)	-0.210* (0.088)	0.255** (0.069)	0.353* (0.163)
Agg. Effect ^a	18582	75690	49223	86510	-38148	-1487	-60815	-53623	645823	939183
Avg. Effect ^b									3.23	4.70
N	890	890	695	695	1084	1084	695	695	2055	2055
Age Trends	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes
Means										
65+, Without Kidney Disease										
Pre	0.96		0.93		0.55		0.04		5.05	
Post	0.98		0.93		0.66		0.04		4.89	
65+, With Kidney Disease										
Pre	0.92		0.91		0.43		0.04		10.35	
Post	0.97		0.93		0.47		0.02		9.43	
<64, Without Kidney Disease										
Pre	0.82		0.00		0.82		0.82		3.87	
Post	0.89		0.04		0.84		0.81		3.89	
<64, With Kidney Disease										
Pre	0.72		0.00		0.76		0.76		9.95	
Post	0.85		0.22		0.64		0.50		11.28	

^a Aggregate effect of the Medicare expansion on insurance coverage and annual number of physician visits for people between 45 and 64 years of age with kidney disease in the post period.

^b Average individual effect of the Medicare expansion on the number of physician visits for people between 45 and 64 years of age with kidney disease in the post period.

Source—Author’s analysis of the National Health Interview Survey from 1968-1980.

Notes—Dependent variable is indicated by the column group title. Kidney disease is defined using the “Broad” definition of kidney disease (see Appendix Table A). DDD is the triple-difference coefficient from the interaction of a dummy for being under 65 years of age, a dummy for the second half of 1973 or later, and a dummy for having kidney disease; DD is the difference-in-difference coefficient from a sample with kidney disease. Models include year, age, gender, and race fixed effects along with all one-, two-, and, if appropriate, three-way interactions of under 65, post, and kidney disease; models with age trends also include additional interactions with age-65. Sample restricted to individuals between 45 and 84 years of age; columns (9) and (10) also exclude people 65 years of age and data from July 1 1973 to June 30 1974. Estimates are from OLS regressions in columns (1)-(8) and Poisson in columns (9) and (10). Standard errors clustered on age in round brackets.

+ p<0.1, * p<0.05, ** p<0.01

Table 3: Poisson Estimates of the Effect of the ESRD Program on Mortality

	Narrow Definition				Broad Definition		Chronic Only	
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
A: Ages 45 to 84								
DDD	-0.373** (0.049)	-0.073* (0.029)	-0.059** (0.007)	-0.021* (0.010)	-0.067** (0.011)	-0.010 (0.009)	-0.057* (0.027)	-0.020+ (0.012)
Δ in Mortality Rate	-2.0	-0.5	-2.3	-0.8	-0.8	-0.5	-0.3	-0.7
Δ in # Deaths	-772	-174	-891	-323	-311	-196	-114	-270
DD	-0.363** (0.053)	-0.079* (0.035)	-0.048** (0.014)	-0.025+ (0.014)	-0.073** (0.022)	-0.015 (0.011)	-0.064+ (0.034)	-0.024+ (0.014)
Δ in Mortality Rate	-2.0	-0.5	-1.9	-1.0	-0.9	-0.7	-0.3	-0.9
Δ in # Deaths	-755	-189	-723	-388	-337	-288	-128	-331
B: Ages 45 to 60 and 70 to 84								
DDD	-0.446** (0.052)	-0.148** (0.049)	-0.080** (0.006)	-0.066** (0.017)	-0.144** (0.021)	-0.040** (0.011)	-0.134* (0.053)	-0.062** (0.021)
DD	-0.437** (0.056)	-0.150* (0.066)	-0.068** (0.015)	-0.065* (0.026)	-0.145** (0.043)	-0.040+ (0.021)	-0.136* (0.069)	-0.062* (0.026)
Age trends	No	Yes	No	Yes	Yes	Yes	Yes	Yes
Underlying only?	Yes	Yes	No	No	Yes	No	Yes	No
Mean Annual Kidney Disease Mortality Rates (per 100,000)								
Ages 45-64								
1968-1973H1	6.6	6.6	40.0	40.0	13.8	52.0	5.6	36.5
1973H2-1978	5.4	5.4	37.3	37.3	8.9	45.3	4.6	32.6
Ages 65-84								
1968-1973H1	28.0	28.0	251.0	251.0	100.4	368.1	24.4	238.2
1973H2-1978	32.4	32.4	241.1	241.1	76.7	323.2	27.4	218.6

Source—Author's analysis of Multiple Cause Mortality Files for 1968-1978.

Notes—Dependent variable is the mortality rate in the age-time-gender-cause of death cells, where time is measured in half-year increments. Definitions of kidney disease based on codes in table A. DDD is the triple difference coefficient for being under 65, in the post expansion period, with kidney disease; DD is the difference-in-differences coefficient for being under 65 and in the post expansion period using a sample that is restricted to deaths due to kidney disease. Change in mortality rate is calculated as the exponentiated coefficient minus 1 multiplied by the pre-period mortality rate; change in number of deaths is the change in the mortality rate multiplied by the population between 45 and 64 years of age in 1973. Models that do not restrict to underlying causes of deaths also define a death as due to kidney disease if kidney disease is either an underlying or a contributing cause of death. All models include time fixed effects (measured in six month increments), age fixed effects, an indicator for female, and all possible interactions of an indicator for being under 65, a post period dummy, and, where appropriate, an indicator for deaths due to kidney disease. Models with age trends also include interactions with age minus 65 in addition to the under 65, post, and kidney disease interactions. Sample is restricted to deaths to whites between 45 and 84 years of age in panels A; panel B excludes deaths to people between 61 and 69 years of age. Estimates are from Poisson regressions, standard errors two-way clustered on age and time in round brackets.

+ $p < 0.1$, * $p < 0.05$, ** $p < 0.01$

Table 4: Cross-Country Estimates of the Effect of the ESRD Program on Kidney Mortality

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
DDDD	-0.064** (0.023)	-0.077** (0.024)	-0.064** (0.024)	-0.076** (0.024)	-0.057** (0.020)	-0.064** (0.023)	-0.058** (0.020)	-0.065** (0.023)
DDD	-0.048* (0.023)	-0.060** (0.022)	-0.045* (0.020)	-0.048* (0.021)	-0.039* (0.017)	-0.040* (0.018)		
Only Members Before 1969	No	Yes	No	Yes	No	Yes	No	Yes
Country FE			X	X	X	X	X	X
Country Interactions					X	X	X	X
Year-by-Kidney							X	X

Source—Authors’ analysis of the World Health Organization Mortality Database for 1968 through 1978, covering the United States and OECD Member States at any point in time.

Notes—Coefficients are point estimates from Poisson regressions using the mortality rate per 100,000 in each country-year-gender-age group-cause of death cell as the dependent variable. DDDD is the coefficient on the four-way interaction of a dummy for the United States, an indicator for the post period, a dummy for deaths due to kidney disease, and a dummy for being 45-64 years of age; DDD is the corresponding coefficient in models that restrict to deaths due to nephritis. All models include year, age, and gender fixed effects and trends in age-65, where age in each cell was recentered by 2.5 years. County Interactions are two-way interactions of country fixed effects with dummies for kidney disease and post. Sample restricted to individuals between the ages of 45 and 84 and years in which the country used the ICD-8 coding regime. Estimates are from Poisson regressions, cells weighted by population, standard errors clustered on country in parentheses.

+ p<0.1, * p<0.05, ** p<0.01, *** p<0.001

Table 5: In-state Treatment Capacity and Mortality Reduction

	Narrow Definition			Chronic Only		
	(1)	(2)	(3)	(4)	(5)	(6)
A: Base Model						
DDD	-0.077** (0.028)	-0.076* (0.031)	-0.078* (0.033)	-0.060* (0.028)	-0.058+ (0.031)	-0.061+ (0.032)
×Log Dialysis Clinics Per Capita in 1971	-0.065** (0.023)		-0.093** (0.033)	-0.062* (0.025)		-0.086* (0.037)
×Log Transplant Programs Per Capita in 1971		-0.008 (0.060)	0.061 (0.063)		-0.009 (0.058)	0.057 (0.065)
B: Including Indicators for VA Treatment Facilities						
DDD	-0.080** (0.029)	-0.077* (0.031)	-0.081* (0.032)	-0.063* (0.029)	-0.059* (0.029)	-0.064* (0.031)
×Log Dialysis Clinics Per Capita in 1971	-0.067** (0.025)		-0.082* (0.037)	-0.065* (0.027)		-0.076* (0.037)
×Log Transplant Programs Per Capita in 1971		-0.016 (0.058)	0.044 (0.065)		-0.017 (0.057)	0.037 (0.064)

Source—Author’s analysis of Multiple Cause Mortality Files for 1968-1978 and the publication “Kidney Disease Services, Facilities, and Programs in the United States” (Kidney Disease Program, 1971).

Notes—Dependent variable is the mortality rate in the state-age-time-gender-cause of death cells, where time is measured in half-year increments. Definitions of kidney disease based on codes in table A. DDD is the triple difference coefficient for being under 65, in the post expansion period, with kidney disease; models with interactions of DDD with either dialysis clinics or transplant programs also include all two- and three- way interactions of dialysis clinics or transplant programs with under 65, post, and the kidney disease indicator. Models also include indicators for having no dialysis clinics or transplant programs in a state; panel B also includes indicators for the presence of VA dialysis clinics and transplant programs (also interacted with DDD). All measures of dialysis clinics and transplant programs have been demeaned. All models include state, time, and age fixed effects, an indicator for female, and age trends interacted with under 65, post, and kidney disease. Sample is restricted to deaths to whites between 45 and 84 years of age. Estimates are from Poisson regressions, standard errors three-way clustered on state, age, and time in round brackets; each state weighted by its total population.
+ p<0.1, * p<0.05, ** p<0.01

Table 6: Entry of Dialysis and Transplant Facilities

	Per 100,000 in 1975					
	Dialysis Clinics			Transplant Programs		
	(1)	(2)	(3)	(4)	(5)	(6)
Log Per Capita						
Dialysis Clinics	0.316*	0.331**	0.360**			
	(0.127)	(0.104)	(0.102)			
Transplant Programs				0.710**	0.617**	0.613**
				(0.154)	(0.129)	(0.135)
Log Kidney Disease Mortality Rate						
Under 65		0.227	0.125		0.370	0.393
		(0.175)	(0.203)		(0.256)	(0.316)
65 and Over			0.320			-0.067
			(0.284)			(0.559)
Constant	-0.381	-0.570*	-1.571	-0.500	-1.140*	-0.936
	(0.260)	(0.257)	(0.963)	(0.437)	(0.572)	(1.782)

Source—Author’s analysis of Multiple Cause Mortality Files for 1971, the publication “Kidney Disease Services, Facilities, and Programs in the United States” (Kidney Disease Program, 1971) and the 1977 Social Security Bulletin. See text for details.

Notes—Independent variables are measured in 1971. Kidney disease mortality defined using the “narrow” definition and underlying causes of death. Estimates from Poisson models, robust standard errors in parentheses.

+ $p < 0.1$, * $p < 0.05$, ** $p < 0.01$

Table 7: Impact of the ESRD Program on Life Expectancy at Age 45

	Narrow Definition				Broad Definition		Chronic Only	
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
Actual								
Survival to age 64	18.36	18.36	18.36	18.36	18.36	18.36	18.36	18.36
Survival to age 84	28.83	28.83	28.83	28.83	28.83	28.83	28.83	28.83
DDD Counterfactual								
Survival to age 64	18.36	18.36	18.36	18.36	18.36	18.36	18.36	18.36
Difference from actual	0.003	0.001	0.003	0.003	0.002	0.002	0.001	0.002
Life years saved	7235	2189	7641	7883	4618	5045	1291	5117
Survival to age 84	28.84	28.83	28.84	28.84	28.84	28.84	28.83	28.84
Difference from actual	0.007	0.002	0.008	0.007	0.005	0.005	0.001	0.005
Life years saved	17671	5085	19546	17451	11025	11201	3043	11655
DD Counterfactual								
Survival to age 64	18.36	18.36	18.36	18.36	18.36	18.36	18.36	18.36
Difference from actual	0.003	0.001	0.003	0.005	0.003	0.005	0.001	0.004
Life years saved	7074	2974	6200	12429	6080	11234	1999	9415
Survival to age 84	28.84	28.83	28.84	28.84	28.84	28.84	28.83	28.84
Difference from actual	0.007	0.003	0.007	0.011	0.006	0.010	0.002	0.009
Life years saved	17278	6699	15857	26992	14152	24274	4500	20687
Age trends	No	Yes	No	Yes	Yes	Yes	Yes	Yes
Underlying only?	Yes	Yes	No	No	Yes	No	Yes	No

Source—Author’s analysis of Multiple Cause Mortality Files for 1968-1978.

Notes—Based on estimates from models presented in table 3. “DDD Counterfactual” uses DDD estimates from table 3 and “DD Counterfactual” uses DD estimates. Counterfactual survival is based on multiplying kidney-specific mortality hazard by the relevant coefficient for cells under age 65 and then computing survival as the sum of the cumulative survival probabilities by age (see text for details). Life years saved based on population of white and non-white 45 year olds.

Tables

Appendix Table A: ICD Codes for Kidney Disease, by ICD Revision

	ICD-7 (1968 NHIS)	ICDA-8 (1968-1978)	ICD-9 (1979-1980 NHIS)
Narrow Definition:			
Chronic Kidney Disease	592-594,792	582-584, 593.2, 792	582-589
Acute Kidney Disease	590-591	580-581, 593.1	580-581, 584
Broad Definition (additions):			
Other Diseases			
of Urinary System	600-609	590-599	590-599
Hypertension	442,446	403-404	403-404
NHIS Omissions	604-609	594-599	594-599

Online Appendix—Not for publication

Online Appendix Figures

A1	Robustness to Different Bandwidths	2
----	--	---

Online Appendix Tables


B1	Effect of the ESRD Program on Health Insurance and Health Care Utilization (Donut Regressions)	3
B2	OLS Estimates of the Effect of the ESRD Program on Mortality . .	4
B3	Robustness of the Effect of the ESRD Program on Mortality	5
B4	The Confounding Effect of the SSI Program on Mortality	6

Figures

Figure A1: Robustness to Different Bandwidths


A: Narrow, Underlying Cause

B: Narrow, Incl. Contributing Cause




C: Broad, Underlying Cause

D: Broad, Incl. Contributing Cause



E: Chronic, Underlying Cause

F: Chronic, Incl. Contributing Cause



Source—Author’s analysis of Multiple Cause of Death Files, 1968-1978.

Notes—See notes to table 3.

Tables

Table B1: Effect of the ESRD Program on Health Insurance and Health Care Utilization (Donut Regressions)

	Any Insurance		Medicare		Any Private		Only Private		Doctor Visits	
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
DDD	-0.037 (0.090)	0.176 (0.169)	0.140* (0.052)	0.455** (0.108)	-0.123 (0.108)	-0.044 (0.229)	-0.277** (0.090)	-0.387* (0.139)	0.170* (0.073)	0.251 (0.246)
Agg. Effect ^a	-7254	34237	27349	88639	-21677	-7793	-53926	-75378	297685	458697
Avg. Effect ^b									2.15	3.32
N	112784	112784	90091	90091	143358	143358	90091	90091	291674	291674
DD	0.020 (0.085)	0.239 (0.194)	0.192* (0.065)	0.500** (0.144)	-0.206* (0.097)	-0.053 (0.272)	-0.260** (0.072)	-0.335* (0.149)	0.236** (0.089)	0.393 (0.254)
Agg. Effect ^a	3803	46626	37363	97435	-36401	-9393	-50728	-65262	462395	836784
Avg. Effect ^b									3.01	5.45
N	636	636	493	493	780	780	493	493	1531	1531
Age Trends	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes

^a Aggregate effect of the Medicare expansion on insurance coverage and annual number of physician visits for people between 45 and 59 years of age with kidney disease in the post period.

^b Average individual effect of the Medicare expansion on the number of physician visits for people between 45 and 59 years of age with kidney disease in the post period.

Source—Author’s analysis of the National Health Interview Survey from 1968-1980.

Notes—Dependent variable is indicated by the column group title. Kidney disease is defined using the “Broad” definition of kidney disease (see Appendix Table A). DDD is the triple-difference coefficient from the interaction of a dummy for being under 65 years of age, a dummy for the second half of 1973 or later, and a dummy for having kidney disease; DD is the difference-in-difference coefficient from a sample with kidney disease. Models include year, age, gender, and race fixed effects along with all one-, two-, and, if appropriate, three-way interactions of under 65, post, and kidney disease; models with age trends also include additional interactions with age-65. Sample restricted to individuals between 45 and 59 and 70 and 84 years of age; columns (9) and (10) also exclude data from July 1 1973 to June 30 1974. Estimates are from OLS regressions in columns (1)-(8) and Poisson in columns (9) and (10). Standard errors clustered on age in round brackets.

+ p<0.1, * p<0.05, ** p<0.01

Table B2: OLS Estimates of the Effect of the ESRD Program on Mortality

	Narrow Definition				Broad Definition		Chronic Only	
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
A: Ages 45 to 84								
DDD	-0.372** (0.052)	-0.060 (0.035)	-0.078** (0.008)	-0.030* (0.012)	-0.068** (0.019)	-0.020* (0.009)	-0.028 (0.033)	-0.028* (0.013)
Δ in Mortality Rate	-2.0	-0.4	-3.0	-1.2	-0.8	-1.0	-0.1	-1.0
Δ in # Deaths	-770	-145	-1162	-456	-313	-398	-58	-377
DD	-0.359** (0.055)	-0.049 (0.041)	-0.063** (0.014)	-0.018 (0.015)	-0.057* (0.023)	-0.009 (0.011)	-0.018 (0.039)	-0.016 (0.015)
Δ in Mortality Rate	-1.9	-0.3	-2.4	-0.7	-0.7	-0.4	-0.1	-0.6
Δ in # Deaths	-748	-119	-942	-275	-263	-172	-36	-219
B: Ages 45 to 60 and 70 to 84								
DDD	-0.453** (0.053)	-0.155** (0.048)	-0.095** (0.009)	-0.079** (0.018)	-0.161** (0.034)	-0.056** (0.012)	-0.105+ (0.053)	-0.073** (0.023)
DD	-0.437** (0.058)	-0.140* (0.064)	-0.077** (0.016)	-0.060* (0.027)	-0.144** (0.048)	-0.038+ (0.021)	-0.090 (0.068)	-0.056+ (0.028)
Age trends	No	Yes	No	Yes	Yes	Yes	Yes	Yes
Underlying only?	Yes	Yes	No	No	Yes	No	Yes	No
Mean Annual Kidney Disease Mortality Rates (per 100,000)								
Ages 45-64								
1968-1973H1	6.6	6.6	40.0	40.0	13.8	52.0	5.6	36.5
1973H2-1978	5.4	5.4	37.3	37.3	8.9	45.3	4.6	32.6
Ages 65-84								
1968-1973H1	28.0	28.0	251.0	251.0	100.4	368.1	24.4	238.2
1973H2-1978	32.4	32.4	241.1	241.1	76.7	323.2	27.4	218.6

Source—Author's analysis of Multiple Cause Mortality Files for 1968-1978.

Notes—Dependent variable is the log mortality rate in the age-time-gender-cause of death cells, where time is measured in half-year increments. Definitions of kidney disease based on codes in table A. DDD is the triple difference coefficient for being under 65, in the post expansion period, with kidney disease; DD is the difference-in-differences coefficient for being under 65 and in the post expansion period using a sample that is restricted to deaths due to kidney disease. Change in mortality rate is calculated as the exponentiated coefficient minus 1 multiplied by the pre-period mortality rate; change in number of deaths is the change in the mortality rate multiplied by the population between 45 and 64 years of age in 1973. Models that do not restrict to underlying causes of deaths also define a death as due to kidney disease if kidney disease is either an underlying or a contributing cause of death. All models include time fixed effects (measured in six month increments), age fixed effects, an indicator for female, and all possible interactions of an indicator for being under 65, a post period dummy, and, where appropriate, an indicator for deaths due to kidney disease. Models with age trends also include interactions with age minus 65 in addition to the under 65, post, and kidney disease interactions. Sample is restricted to deaths to whites between 45 and 84 years of age in panels A; panel B excludes deaths to people between 61 and 69 years of age. Estimates are from OLS regressions, standard errors two-way clustered on age and time in round brackets.

+ $p < 0.1$, * $p < 0.05$, ** $p < 0.01$

Table B3: Robustness of the Effect of the ESRD Program on Mortality

Power	Underlying Cause of Death					Underlying and Contributing Cause of Death				
	Age Trends		Time Trends			Age Trends		Time Trends		
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
N/A	Linear	Quadratic	Linear	Quadratic	N/A	Linear	Quadratic	Linear	Quadratic	
A: Narrow Definition										
DDD	-0.373** (0.049)	-0.073* (0.029)	-0.059* (0.029)	-0.076 (0.056)	-0.075 (0.066)	-0.059** (0.007)	-0.021* (0.010)	-0.016 (0.010)	-0.017* (0.007)	-0.020** (0.007)
DD	-0.363** (0.053)	-0.079* (0.035)	-0.064* (0.031)	-0.067 (0.062)	-0.062 (0.071)	-0.048** (0.014)	-0.025+ (0.014)	-0.018 (0.013)	-0.009 (0.018)	-0.007 (0.018)
B: Broad Definition										
DDD	-0.197** (0.019)	-0.067** (0.011)	-0.048** (0.012)	-0.073** (0.010)	-0.070** (0.013)	-0.042** (0.006)	-0.010 (0.009)	-0.007 (0.008)	-0.030** (0.006)	-0.030** (0.007)
DD	-0.189** (0.025)	-0.073** (0.022)	-0.053** (0.017)	-0.066* (0.028)	-0.058+ (0.033)	-0.032** (0.010)	-0.015 (0.011)	-0.010 (0.010)	-0.021 (0.015)	-0.017 (0.014)
C: Chronic Definition										
DDD	-0.340** (0.046)	-0.057* (0.027)	-0.042 (0.026)	-0.062 (0.058)	-0.063 (0.067)	-0.056** (0.007)	-0.020+ (0.012)	-0.014 (0.011)	-0.018* (0.007)	-0.022** (0.007)
DD	-0.331** (0.051)	-0.064+ (0.034)	-0.047 (0.029)	-0.053 (0.064)	-0.050 (0.074)	-0.045** (0.013)	-0.024+ (0.014)	-0.017 (0.012)	-0.010 (0.019)	-0.009 (0.020)

Source—Author’s analysis of Multiple Cause Mortality Files for 1968-1978.

Notes—See Note to table 3.

+ p<0.1, * p<0.05, ** p<0.01

Table B4: The Confounding Effect of the SSI Program on Mortality

	Narrow Definition		Broad Definition		Chronic Only	
	(1)	(2)	(3)	(4)	(5)	(6)
DDD	-0.0572 (0.039)	-0.0814** (0.030)	-0.0654* (0.030)	-0.0809** (0.017)	-0.0356 (0.035)	-0.0562+ (0.029)
×% with Soc. Security Income	0.114 (0.125)		0.0145 (0.081)		0.0509 (0.130)	
×% with SSI Income	-0.00173 (0.783)		-0.580 (0.563)		0.447 (0.769)	
×More Stringent Criteria		0.0491 (0.039)		0.0357+ (0.019)		0.0327 (0.040)
×Separate Medicaid Application		-0.0124 (0.035)		0.0440* (0.019)		-0.0279 (0.051)

Source—Author’s analysis of Multiple Cause Mortality Files for 1968-1978 and the Current Population Survey.

Notes—Dependent variable is the mortality rate in the state-age-time-gender-cause of death cells, where time is measured in half-year increments. Definitions of kidney disease based on codes in table A. DDD is the triple difference coefficient for being under 65, in the post expansion period, with kidney disease. % with Soc. Security Income and % with SSI income are the demeaned percentages of people in a state-age-gender cell who report receiving either Social Security or SSI income, respectively, in the 1977 to 1979 March CPS. More Stringent Criteria indicates that a state has adopted a more stringent asset or income test for Medicaid eligibility than used by the SSI program, and Separate Medicaid Application indicates that a state requires individuals to separately apply for SSI and Medicaid. Interactions with DDD also include interactions with under 65, post, and kidney disease. All models include state, time, and age fixed effects, an indicator for female, and age trends interacted with under 65, post, and kidney disease. Sample is restricted to deaths to whites between 45 and 84 years of age. Estimates are from Poisson regressions, standard errors three-way clustered on state, age, and time in round brackets; each state weighted by its total population.

+ $p < 0.1$, * $p < 0.05$, ** $p < 0.01$